



RESEARCH ARTICLE

Pathogenicity of *Colletotrichum* species isolates on different mango (*Mangifera indica* L.) varieties in Southwestern Ethiopia

Amsalu Abera¹, Fikre Lemessa², Girma Adunga², Gerba Daba²

¹ Ambo University, College of Agriculture and Veterinary Sciences, Department of Plant Sciences, P. O. Box 19 Ambo, Ethiopia,

² Jimma University College of Agriculture and Veterinary Medicine, Department of Horticulture and Plant Sciences, P.O. Box 307, Jimma, Ethiopia

Received: 26.08.2022

Accepted: 08.10.2022

ABSTRACT

Mango (*Mangifera indica* L.) is one of the most popular fruits to millions of people and grown throughout the tropical and subtropical regions of the world. *Colletotrichum* species are the most important and prevalent pathogen causing great yield losses to mango growers in the country. Characterization of the causal agent by pathogenicity test and virulence variability of the pathogen on their host response is not well documented in southwestern part of Ethiopia. Therefore, the aims of the current study were to investigate pathogenicity of *Colletotrichum* species (isolates) and to assess pathogenic variability on local and commercially cultivated mango varieties in Ethiopia. Isolates of *Colletotrichum* species collected from nine locations in southwestern Ethiopia from mango leaves, panicles and fruit were studied for pathogenicity. Artificial inoculation of 5 varieties of Tommy Atkins, Keitt, Kent, Apple mango, Vandyke and local variety was made by wounding detached mango fruits, leaves and seedlings under greenhouse condition. The pathogenicity test result on detached mango fruits showed the presence of three virulence level among *Colletotrichum* isolates as highly pathogenic (62.5%), pathogenic (25%) and mildly pathogenic (12.5%). Virulent isolate tested on mango seedlings scored (13.6mm) lesion diameter. Accordingly, local mango varieties, Kent and Vandyke were categorized as susceptible varieties, while Tommy Atkins, Keitt and Apple mango varieties were intermediately susceptible. Since this result was solely based on laboratory experiment, we recommend further research to be conducted at field conditions on those varieties and including other mango varieties in the country.

Keywords: *Colletotrichum*, Ethiopia, Keitt, Kent, mango, pathogenicity, postharvest

Citation: Abera, A., Lemessa, F., Adunga, G., and Daba, G. 2022. Pathogenicity of *Colletotrichum* species isolates on different mango (*Mangifera indica* L.) varieties in Southwestern Ethiopia. *Journal of Postharvest Technology*, 10(4): 113-123.

INTRODUCTION

Mango (*Mangifera indica* L.) is ranked as one of the best fruits in the international market because of its delicious taste and high caloric value. This fruit has become an essential fruit crop in Asia, Southern and Central America as well as in many parts of Africa including Ethiopia (Phoulivong et al., 2010a).

* For correspondence: A. Abera (Email: aamsalu97@gmail.com)

Various biotic and abiotic stresses are causing immense loss to mango crop throughout the world. Among biotic stresses, anthracnose of mango, caused by *Colletotrichum* species is the most serious and wide spread disease throughout the mango production areas, leading to substantial pre and post-harvest losses of the crop (Syed et al., 2014). The pathogen causes irregularly shaped and black necrotic spots of symptoms on both sides of mango leaves. In the inflorescences, the symptoms appear in the stalk and in individual flowers. Although the fungus affects many parts of the plant, the most significant losses occur in the ripening and post-harvest stages, when black, sunken and necrotic lesions would appear on the peel of the mango fruit (Meer et al., 2013). There is large variation among and within *Colletotrichum* species in pathogenicity, culture appearance, symptom and uncertain relationships with host plants (Sutton, 1992).

The symptom expression varies among isolates, for the resistant and susceptible mango varieties. The resistant level of different genotypes against anthracnose in different crops such as chili pepper have been screened (Prasath et al., 2007), grapevine (Shelke et al., 1997) and mango (Pandey et al., 2011). Host-pathogen interactions depended on expression of resistance factors by the mango varieties and pathogenicity factors by the fungus during different periods of fruit life (Quang and Somsiri, 2002).

The economic importance of anthracnose on mango in Ethiopia commands the need for studies on the identification and evaluation of the pathogenicity of *Colletotrichum* isolates on their host. However, characterization of the causal agent by pathogenicity test and virulence variability of the pathogen on their host response is not well documented in southwestern part of Ethiopia. This study was initiated to investigate the pathogenicity of *Colletotrichum* species isolates associated with mango and pathogenic variability on existing and commercially cultivated mango varieties in Ethiopia.

MATERIALS AND METHODS

Colletotrichum species isolates were retrieved from the leaves, panicles and attached fruit of mango showing typical symptoms of anthracnose. The isolates were taken from nine location of mango grown at home gardens southwest Ethiopia. After which the isolates were grown on potato dextrose agar medium (PDA). After confirming the spores, the cultures were purified by single spore isolation technique (Dhingra and Sinclair, 1993). The fungi were identified on the basis of morphological characteristics through mycological key suggested by Barnett et al. (1972). Suspension of conidia was prepared by suspending mycelia scraped from seven days old of *Colletotrichum* species isolates were separately added in 3-milliliter sterile distilled water and shaken vigorously for 3 minutes. The resulting suspensions were filtered through 2-layer of cheese cloth. The concentrations of spore suspension were adjusted to (1×10^6 spore's ml⁻¹) or conidia using haemocytometer.

Artificial inoculation methods were commonly used to test the pathogenicity of a fungal species *in vitro*, as it is easy to control environmental conditions (Ekbote et al., 1997). Pathogenicity tests on detached fruits, leaves and seedling of mango varieties (cv. Tommy Atkins, Keitt, Kent, Apple mango, Vandyke and locale mango) collected from Melkasa Agricultural Research Center was made. Detached fruits and leaves were washed under running tap water for 60 seconds followed by surface sterilization by immersing the fruits in 70% ethanol for 3 minutes, then rinsed three times in sterile distilled water for 2 minutes and dried with sterile tissue paper and then air dried.

Inoculations of isolates on physiologically matured detached mango fruits were done. , Surface sterilized fruits were placed in a plastic box having three fruits per treatment with tissue paper then sprayed with sterilized water to maintain at least 95% relative humidity (Than *et al.*, 2008a). The samples were inoculated using the wound inoculation method which included pin-pricking the fruits to a 3 mm depth with a sterile needle in the middle portion of fruit, and then placing 20 μ l of conidia suspension

on to the wound (Than et al., 2008a, b). Control fruits were inoculated with 20µl of sterile distilled water. The inoculated samples were incubated in the plastic containers at 25°C under controlled conditions. The plastic box was removed after 48hrs and fruits were kept at the same temperature. Fruits were monitored for the onset of symptoms for 5-11 days. Pathogenicity test on detached leaves free from anthracnose symptom were collected from greenhouse grown seedlings of Vandyke, Keitt, Tommy Atkins, Kent, Apple mango and local mango varieties. The leaves were properly washed, and surface sterilized before treatment. Then three leaves per treatments were placed on plastic boxes lined on the inside with moist tissue paper. Wound inoculations technique was performed following the protocol described by Oliveira Costa et al. (2010). Aqueous conidial suspensions of *Colletotrichum* isolates (1×10^6 spore's ml⁻¹) were prepared from 7-day-old cultures of each isolates. Leaves inoculated with sterile distilled water drops served as controls. Inoculated leaves were incubated in plastic boxes with moist tissue paper for 48 hours in the laboratory to ensure high humidity that creates favorable conditions for conidial germination and infection at room temperature. The inoculated leaves were monitored daily after 5 days of inoculation for lesions formation and measured accordingly.

Anthraco development on seedling test, six mango varieties were arranged in the greenhouse and three attached leaves of all varieties having similar size were selected for inoculation. The selected leaves were washed, and surface sterilized by immersing the leaves in 70% ethanol for 3 minutes and then immersing three times in sterile distilled water for 2 minutes and drying with sterile tissue paper and then air dried (Sanders and Korsten, 2003). Then 20µl of conidia suspension was placed onto three spot on each leave, and the control tissues was similarly treated with distilled water. Inoculated seedling leaves were incubated in moist plastic sheet for 48 hours in greenhouse condition to ensure high humidity for establishing favorable conditions for conidial germination and infection.

Disease reactions of the hosts were evaluated by measuring the length, width and area of the typical anthracnose lesion which developed on tissues. Isolates were considered pathogenic when the lesion area advanced beyond the 4mm diameter initial injury. The pathogenicity of *Colletotrichum* strains was estimated according to the grading scale (Nova et al., 2011) presented in the (Table 1) varying from 0 to 4. Lesion length (along the long axis of the tissues) and width (along the short axis of the tissues) were measured and evaluation of response of different mango varieties against anthracnose disease of mango seedling caused by *Colletotrichum* isolate at (Table .2) greenhouse (Giblin et al., 2010). It was assumed that lesions grew in a circular manner so that their diameter (LD) was evaluated using the formula proposed by Sharma et al. (2005).

$$LD \text{ (mm)} = \frac{\text{Lesion length} + \text{lesion width}}{2}$$

The virulence of the isolates was evaluated by measuring the lesion length at 11 days after inoculation in two perpendicular directions on each leaves. Lesion diameter and Percent Disease Index (PDI) was calculated. The PDI was calculated by using the formula (Prabakar et al., 2005).

$$PDI = \frac{\text{Sum of all numerical ratings}}{\text{Total number of observations}} \times \frac{100}{\text{Maximum rating observed}}$$

The numerical values of percent disease index and latent period were used to calculate the virulence index using the following formula (Thakur and Rao, 1997).

Virulence index (VI) = Percent disease index (PDI) × Latent period⁻¹

Table 1: Grading scale used for evaluation of the pathogenicity of *Colletotrichum* species isolates in mango fruits (Nova et al., 2011)

Grade	Symptoms characteristics (lesions mm)	Classification of the isolates
0	Absence of lesions	Non-pathogenic
1	from 1 to 5 mm	Weakly pathogenic
2	Lesions from 6 to 15 mm	Mildly pathogenic
3	from 16 to 25 mm	Pathogenic
4	Lesions over 25 mm	Highly pathogenic

Table 2: Grading scale used for evaluation of response of different mango varieties against anthracnose disease of mango seedling caused by *Colletotrichum* isolate at greenhouse (Giblin et al., 2010)

Scale rate	Symptom lesion(mm)	Host response
0	no symptom	Immune
1	1mm lesion	Highly resistance
2	2-3mm lesion	resistance
3	3-5mm lesion	intermediate
4	lesion>5mm	susceptible
5	spread all area of leaves	highly susceptible

Statistical analysis

The laboratory and greenhouse experiments were arranged in Completely Randomized Design (CRD) with three replications. The experiment was carried out twice for each experiment. Single and interaction effects of factors were determined using the general liner model (GLM) procedure of (SAS, 2008). Whenever significant interaction was observed between factors; the level of one factor was compared at level of other factors. Mean values among treatments were compared by the LSD at p= 5% level of significance.

RESULTS AND DISCUSSION

The finding revealed that the interaction between *Colletotrichum* species isolates and tested mango varieties showed significant effect on the pathogenicity test of the physiologically matured detached mango fruits ($p < 0.0001$). Among eight isolates of *Colletotrichum* tested, maximum disease severity index 30.5, 33.3, 30.5, 30.4, were recorded in CGm-F, CSC, CBK-F, CAD-F and CBL-L, respectively (Table 3) on Vandyke mango variety while the severity index was not significantly different from each other. The result indicated that lower severity (percent disease index), was observed on isolate CMn-w (11.1 %) on Apple mango (Table 3). According to their pathogenicity, *Colletotrichum* isolates based on mean lesion diameters were grouped into three as highly pathogenic, pathogenic and mildly pathogenic (Table 4).

Table 3: Disease severities index (%) of *Colletotrichum* isolates on detached physiological matured mango fruits

Isolates	Varieties					
	Vandyke	Keitt	Tommy Atkins	Kent	Apple mango	Local mango
CGm-F	30.5 ^{ab}	33.3 ^a	33.1 ^a	33.2 ^a	32.8 ^a	33.3 ^a
CAd-F	30.4 ^{ab}	33.1 ^a	33.2 ^a	33.1 ^a	33.1 ^a	33.0 ^a
CBk-F	30.5 ^{ab}	33.2 ^a	27.7 ^{bc}	33.3 ^a	25.2 ^{cd}	33.2 ^a
CMn-w	25.3 ^{cd}	25.2 ^{cd}	16.7 ^f	25.1 ^{cd}	11.1 ^g	30.5 ^{ab}
CBm-F	20.5 ^{cd}	16.7 ^f	16.6 ^f	27.2 ^{bc}	30.1 ^{ab}	16.7 ^f
CG-L	19.4 ^{ef}	25.0 ^{cd}	22.2 ^{de}	22.1 ^{de}	27.7 ^{bc}	19.4 ^{ef}
CSC	33.3 ^a	33.1 ^a	33.0 ^a	33.3 ^a	33.3 ^a	33.0 ^a
CBL-L	30.5 ^{ab}	33.2 ^a	27.8 ^{bc}	30.1 ^{ab}	27.4 ^{bc}	29.1 ^{ab}
Control	0.0 ^h	0.0 ^h	0.0 ^h	0.0 ^h	0.0 ^h	0.0 ^h

Means followed by the same letter are not significantly different at p= 5% level from each other. LSD values for the interactions comparisons were: 4.8 and CV=10.4%.

Isolates CGm-F, CAd-F, CBk-F, CSC and CBL-L were highly pathogenic, but isolates CGL and CMn-w were pathogenic, while isolate CBm-F was mildly pathogenic except on Kent variety (Table 4).

Table 4: Pathogenicity of *Colletotrichum* isolates on detached physiological matured mango fruit lesion diameter (mm)

Isolates	Varieties					
	Vandyke	Keitt	Tommy Atkins	Kent	Apple mango	Local mango
CGm-F	25.3 ^{h-l}	45.3 ^{abc}	35.6 ^f	42.0 ^{cde}	41.6 ^{cde}	47.6 ^{ab}
CAd-F	24.0 ^{i-l}	38.3 ^{d-f}	30.0 ^{gh}	38.0 ^{def}	34.3 ^{fg}	41.6 ^{cde}
CBk-F	26.0 ^{h-j}	34.0 ^{fg}	26.0 ^{hij}	34.7 ^{fg}	23.0 ^{h-l}	35.0 ^{fg}
CMn-w	18.0 ^{m-p}	21.0 ^{j-n}	12.6 ^{q-s}	20.3 ^{k-n}	20.0 ^{l-o}	25.3 ^{h-l}
CBm-F	20.3 ^{k-n}	10.0 ^{rs}	11.5 ^{p-s}	24.6 ^{h-l}	6.0 ^s	10.3 ^{rs}
CGL	14.7 ^{o-r}	22.3 ^{i-l}	16.0 ^{q-r}	14.7 ^{o-r}	25.7 ^{h-k}	14.3 ^{o-r}
CSC	29.7 ^{gh}	36.7 ^{fg}	33.7 ^{fg}	30.0 ^{gh}	50.0 ^a	43.3 ^{bcd}
CBL-L	27.0 ^{hi}	46.0 ^{abc}	26.3 ^{hij}	35.0 ^{fg}	47.7 ^{ab}	41.3 ^{cde}
Control	0.0 ^t	0.0 ^t	0.0 ^t	0.0 ^t	0.0 ^t	0.0 ^t

Means followed by the same letter are not significantly different at p= 5% level from each other. LSD values for the interactions comparisons were 5.5 and CV=12.1%.

The pathogenicity test result on detached mango fruits showed that the virulence variability of *Colletotrichum* isolates, the result was in line by Pandey et al. (2011) who reported *C. gloeosporioides* isolates produced the highest lesion growth compared to other isolates. It can be argued that variation in the isolates may be inherent since the isolates were collected from different sites. Hence, the physiological characters are influenced by environmental conditions through natural mutations which may be

responsible for such variability in virulence. Similar results were obtained by Rojas-Martinez et al. (2008), who have showed differences in the virulence of the *C. gloeosporioides* isolates.

The virulent isolate CGm-F was used to test for varietal susceptibility on six different mango varieties namely Van-Dyke, Keitt, Tommy Atkins, Kent, Apple mango, and Local mango. The results revealed that, the range of lesion diameter on 5 DAI was from 8.4 to 14.4 mm (Table 5). Typical anthracnose lesions were produced by CGm-F isolate on all the varieties tested. The development of anthracnose symptoms on different varieties were statistically significant ($p < 0.001$) compared to lesion diameter at 5, 7, 9 and 11 days after inoculation on the fruits.

The size of the lesions was gradually increased as day's progress. Eleven days after the inoculation, the lesion size was significantly higher in local mango (32.3 mm) followed by Keitt (31.7 mm) and Kent (29.9mm), but there was no significant difference between them. The least lesion size was recorded in Tommy Atkins (23.9 mm), on 11 days after inoculation (Table 5). The present finding suggests that the *Colletotrichum* isolates were more pathogenic on detached fruits than seedlings leaves. This might be due to accessibility of substrates in fruits (Prusky and Plumbley, 1992).The authors further notes that the susceptibility of fruits to *Colletotrichum* infection is related to the level of antifungal inhibitors present in these fruits. At maturity, hydrolytic process is triggered and poses the depletion of starch, and degradation of insoluble pectin (Phan, 1987).

Table 5: Reactions of *Colletotrichum* isolate CGm-F on detached mango fruits of different mango varieties expressed as diameter of lesion progress days after isolate inoculation (DAI).

Varieties	Disease reaction			
	Diameter of lesion(mm) Days after inoculation (DAI)			
	5DAI	7DAI	9DAI	11DAI
Vandyke	8.4 ^d	15.6 ^d	18.6 ^d	29.1 ^{ba}
Keitt	11.9 ^b	22.0 ^a	26.6 ^a	31.7 ^{ba}
Tommy-Atkins	10.7 ^c	18.5 ^{bc}	21.5 ^c	23.9 ^c
Kent	12.0 ^b	19.3 ^b	23.4 ^{cb}	29.9 ^{ba}
Apple Mango	14.4 ^a	20.2 ^{ba}	23.0 ^{cb}	28.2 ^b
Local Mango	11.8 ^{cb}	17.2 ^{dc}	24.0 ^b	32.3 ^a
Control	0.0 ^e	0.0 ^e	0.0 ^e	0.0 ^d
LSD 5%	1.1	1.9	1.9	3.4
CV (%)	19.9	21.5	17	21.2

*In a column, means followed by the same letters are not significantly different at $p = 5\%$ level

Among the six different mango varieties tested, the lesion diameter was constantly increasing in local mango variety compared to others over the incubation period. In addition, it was considered susceptible to the virulent isolates and showed statistically significant result ($P < 0.001$).

Pathogenicity tests on detached mango leaves, eleven days' after inoculation, all the isolates caused lesions (6.2-12.6 mm) on local mango variety detached leaves with statistically significant ($P < 0.001$) among them. On detached leaves the most aggressive isolate was CBL-L and CGm-F with a mean lesion diameter of 12.6 and 11.2 mm, followed by CAd-F with a mean lesion diameter of 10.7 mm respectively (Table 6). However, other isolates produced lesions (6.2-9.8 mm).

Table 6: Virulence variability in *Colletotrichum* isolates on detached on local mango leaves

<i>Colletotrichum</i> isolates	Disease		
	Lesion diameter (mm)	Percent of disease index	Virulence index
CGm-F	11.2 ^b	25.7 ^{ba}	2.7 ^{ba}
CAd-F	10.7 ^{bc}	24.1 ^{bc}	2.5 ^{bc}
CBk-F	9.8 ^c	23.1 ^c	2.4 ^c
CMn-w	6.8 ^d	19.6 ^d	2.0 ^d
CBm-F	6.7 ^d	20.3 ^d	2.1 ^d
CGL	6.2 ^d	18.4 ^d	1.9 ^d
CSC	9.3 ^c	22.8 ^c	2.4 ^c
CBL-L	12.6 ^a	26.6 ^a	2.8 ^a
Control	0.0 ^e	0.0 ^e	0.0 ^e
LSD 5%	1.35	2.4	0.2
CV (%)	22.2	14.3	14.35

* In a column, means followed by a common letter are not significantly different at p = 0.05 level

All evaluated isolates caused typical anthracnose symptoms on the detached local mango leaves. The leaves that served as controls did not exhibit any symptoms. Some isolates were able to induce extensive lesions on the leaves at 100% of inoculated points. However, in some isolates the infection was only 5% of the inoculated points. Isolates (CGm-F, CAd-F and CBk-F) caused rounded black spots, whereas the necrotic lesions caused by isolates (CMn-w, CBm-F and CGL) and isolates (CSC, and CBL-L) caused lesion surrounded by necrotic and black spots, respectively, indicating the region of growth of the fungus in the leaf tissue. Evaluation of virulence of isolates on mango seedling/attached leaves. The result of the study revealed that disease severity index was significantly varied and the disease severity ranged between 6.7–33.4% (Table 7). Disease severity and lesion developments interaction effect between isolates and mango varieties showed significant difference ($P < 0.0001$) at greenhouse.

Table 7: Disease severity (%) of *Colletotrichum* isolates on 2 years' different varieties of mango seedling/leaves

Isolates	Varieties					
	Vandyke	Keitt	Tommy Atkins	Kent	Apple mango	Local mango
CGm-F	24.6 ^{b-d}	17.7 ^{d-g}	13.3 ^{fi}	33.3 ^a	28.8 ^{ab}	33.4 ^a
CAd-F	20.0 ^{c-f}	13.3 ^{fi}	6.7 ⁱ	17.8 ^{d-g}	13.3 ^{fi}	15.5 ^{e-h}
CBk-F	15.5 ^{e-h}	20.0 ^{c-f}	13.3 ^{fi}	22.2 ^{b-e}	8.9 ^{hi}	24.5 ^{b-d}
CMn-w	15.5 ^{e-h}	24.4 ^{b-d}	13.3 ^{fi}	13.3 ^{fi}	13.3 ^{fi}	28.8 ^{ab}
CBm-F	17.7 ^{d-g}	17.7 ^{d-g}	11.1 ^{g-i}	33.3 ^a	11.1 ^{g-i}	26.6 ^{a-c}
CG-L	22.2 ^{b-e}	11.1 ^{g-i}	6.7 ⁱ	24.4 ^{b-d}	13.0 ^{fi}	15.5 ^{e-h}
CSC	22.1 ^{b-e}	13.3 ^{fi}	13.3 ^{fi}	22.2 ^{b-e}	13.3 ^{fi}	22.2 ^{b-e}
CBL-L	22.3 ^{b-e}	13.3 ^{fi}	17.6 ^{d-g}	20.0 ^{c-f}	33.3 ^a	33.3 ^a
Control	0.0 ^j	0.0 ^j	0.0 ^j	0.0 ^j	0.0 ^j	0.0 ^j

In a column, means followed by the same letter are not significantly different at p= 5% level from each other. LSD values for the interactions comparisons were 7.4, and CV=24.5%.

The isolates CGm-F, CMn-w and CBm-F (33.4mm, 28.8mm and 26.6mm) showed the highest disease severity index on mango seedling respectively but the least severity index were recorded (6.7mm and 11.1mm) by isolates CG-L and CBm-T on varieties Tommy Atkins, Apple mango and Keitt.

Table 8: Response of different mango seedling against anthracnose disease caused by *Colletotrichum* isolates under greenhouse condition lesion diameter (mm) on leaves.

Verities	<i>Colletotrichum</i> isolates							
	CGm-F	CAd-F	CBk-F	CMn-w	CBm-T	CG-L	CSC	CBL-L
Vandyke	9.1 ^b	5.9 ^{h-l}	4 ^{n-s}	5.2 ⁱ⁻ⁿ	6.5 ^{e-h}	7.0 ^{d-g}	8.2 ^{b-d}	5.3 ^{h-m}
Keitt	6.8 ^{e-g}	4.0 ^{k-o}	3.3 ^{p-u}	4.9 ^{j-o}	4.1 ^{k-o}	3.2 ^{p-u}	3.3 ^{p-u}	3.3 ^{p-u}
Tommy-A	4.4 ^{m-q}	3 ^{q-v}	3.0 ^{q-u}	4.3 ^{m-q}	2.6 ^{t-v}	2.0 ^v	5.2 ^{i-o}	7.3 ^{c-f}
Kent	13.6 ^a	3.5 ^{p-v}	5.0 ^{t-o}	3.9 ^{q-s}	6.0 ^{g-k}	5.1 ^{i-o}	4.9 ^{j-o}	4.4 ^{m-q}
Apple M	7.4 ^{c-f}	2.7 ^{s-v}	3.0 ^{q-v}	3.3 ^{p-u}	2.9 ^{s-v}	2.4 ^{uv}	4.1 ^{m-r}	6.3 ^{ei}
Local M	6.0 ^{g-k}	4.6 ^{l-p}	5.3 ^{h-m}	6.2 ^{f-j}	8.3 ^{bc}	4.9 ^{j-o}	5.1 ^{j-o}	8.3 ^{bc}

In a column, means followed by the same letter are not significantly different at p= 5% level from each other. LSD values for the interactions comparisons were 1.33, and CV=16%.

Greenhouse screening of five commercials and one local varieties of mango were performed in pots for the identification of resistance potential against anthracnose of mango caused by *Colletotrichum* isolates. Progress of the disease was measured in diameter of lesion (mm) along the inoculated leaves. The interaction effect between isolates and mango varieties showed significant difference ($P < 0.0001$). Maximum mean lesion diameter 13.6 mm, 9.1 mm and 8.3mm was developed on Kent, Vandyke and local mango varieties after 11 days of inoculation respectively. Minimum mean lesion diameter was exhibited by Tommy Atkins (2mm) and Apple mango (2.4mm) respectively (Table 8). The result showed that most isolates, to varying degrees produced anthracnose symptoms on mango tissues. Isolates of *Colletotrichum* inoculated to detached mango fruit, in the laboratory were more aggressive on mango inoculated on detached leaves and mango seedling. This might be true because the leaf and fruit are very much different in physical structure and chemical composition. Each isolate of the pathogen studied was found to induce anthracnose on the different varieties of mangos. The findings are in agreement with the results of several authors that revealed *Colletotrichum* species are the causal agent of mango (Onyeani et al., 2012).

Varieties of mango seedling inoculated with *Colletotrichum* isolates in the greenhouse were infected and exhibited lesions typical of those observed in the field. Lesions began as dark brown, sunken, circular to irregular spots with distinctly gray centers. Re-isolation of *Colletotrichum* isolate CGm-F from inoculated seedling was consistent and confirmed the pathogenic role. According to the present results it is an intermediately susceptible to the pathogen. The result confirms the finding of Jayasinghe and Fernando (2009) who have report that most or all commercial mango varieties are susceptible to mango anthracnose, some are less susceptible than others, the variety Keitt, less susceptible than Kent, while Kensington Pride, the widely grown cultivar in Australia, is listed as moderately resistant to the disease.

CONCLUSION

Among eight isolates used, CGm-F was identified as virulent isolate based on lesion diameter, percent disease index and virulence index. Among six mango varieties, local mango and Kent were susceptible varieties. None of mango varieties namely

Keitt, Kent, Vandyke, Apple mango, local mango and Tommy Atkins were resistant to the mango anthracnose disease in Ethiopia. Varieties of Tommy Atkins, Keitt and Apple mango were intermediately susceptible varieties. Varieties that have shown intermediate susceptibility in the present experiment may further be tested under different agro-climatic regions of mango producing areas.

ACKNOWLEDGMENT

We would like to thank Jimma University College of Agriculture and veterinary medicine for funding for this work.

REFERENCES


- Barnett, H.L., and Hunter B.B. 1972. Illustrated Genera of Imperfect Fungi. Burgess publishing company: Minneapolis MN, USA.
- Dhingra, O. D. and Sinclair, J.B., 1993. Basic Plant Pathology methods. CBS Publications and Distribution, New Delhi, India. p. 335.
- Ekbote, S.D., Padaganur, G.M., Patil, M.S. and Chattannavar, S.N., 1997. Studies on the cultural and nutritional aspects of *Colletotrichum gloeosporioides*, the causal organism of mango anthracnose. *Journal of Mycology Plant Pathology*, 27: 229–230.
- Giblin, F.R. L.M Coasts and J.A.G. Irwin, 2010. Pathogenic diversity of avocado and mango isolates of *Colletotrichum gloeosporioides* causing anthracnose and paper spot in Australia. *Australas Plant Pathology*, 39:50-62.
- Jayasinghe, C.K., Fernando T., 2009. First report of *Colletotrichum acutatum* on *Mangifera indica* in Sri Lanka. *Ceylon Journal of Science*, 38:31–4.
- Lazan, H., Z.M.A. Ali, K.W. Lee, J. Voon and G.R. Chaplin., 1986. The potential role of polygalacturonase in pectin degradation and softening in mango. *Asean Food Journal*, 2: 93-95.
- Lima, N.B, Batista M.V.A, De Morais J.M.A, Barbosa M.A.G, Michereff S.J, Hyde K.D, Camara M.P. S, 2013. Five *Colletotrichum* species are responsible for mango anthracnose in north eastern Brazil. *Fungal Diversity*, 61:75-88.
- Martinez, E., Osorio J.A., Torres M.F., 2009. Identification of *Colletotrichum* species causing anthracnose on Tahiti lime, tree tomato and mango. *Agronomía Colombiana*, 27: 211–18.
- Meer, H., Shazialram, Iftikhar Ahmad, Faisal SohailFateh, MunawarRazaKazmi, 2013. Identification and Characterization of Post-Harvest Fungal Pathogens of Mango from Domestic Markets of Punjab. *Intenational Journal of Plant Production*, 4: 650-658.
- Nova, V., M. X. V, Borges L. R, De Sovsa, Brasileiro E.A. Lima L.A da Costa A.F de Oliveria, 2011. Pathogenicity for Onion and genetic diversity of isolates of pathogenic fungi *Colletotrichum gloeosporioides* (phyllochoraceae) from the state of pernanmbuses. *Birazil. Genetetics and Molecular Research*, 10:311-320

- Oliveira Costa V.S., Michereff S.J., Martins R.B., Gava C.A. T., Mizubuti E.S.G. and Câmara M.P.S., 2010. Species of Botryosphaeriaceae associated on mango in Brazil. *European Journal of Plant Pathology*, 127:509–519.
- Onyeani, C.A Osunlaja, S. Oworu O. O and Olufemi S., 2012. First Report of Fruit Anthracnose in Mango caused by *Colletotrichum gloeosporioides* in Southwestern Nigeria. *International Journal of Agricultural Research*, 4:30-34
- Pandey, A., Pandey, B.K., Muthukumar, M., Rajan, S., Yadava, L.P. and Chauhan, U.K. 2011. In vitro screening for resistance to anthracnose (*Colletotrichum gloeosporioides*) in different mango (*Mangifera indica*) hybrids. *Tunis. Journal of Plant Protection*, 6:99- 108.
- Phan, C.T., 1987. Biochemical and physiological changes during the harvest period, pp. 9-21. In J. Weichmann, ed. *Postharvest physiology of vegetable*. Marcel Dekker Inc., New York
- Phoulivong, S., Cai L., Chen H., McKenzie E.H.C., Abdelsalam K.E.C, Hyde K.D, 2010a. *Colletotrichum gloeosporioides* is not a common pathogen on tropical fruits. *Fungal Diversity*, 44: 33–43.
- Prabakar, K., Raguchander, T., Parthiban, V.K., Muthulakshmi, P., and Prakasam, V., 2005. Post harvest diseases and disorders of mango and their management. In: S.A.M.H. Naqvi (ed.). *Diseases of Fruits and Vegetables*, pp 511 – 619, 2004.
- Prasath, D., Ponnuswami, V., and Muralidharan, V., 2007. Source of resistance to anthracnose (*Colletotrichum capsici*) disease in *Capsicum* species. *Indian Journal of Agricultural Research*, 77: 473-474.
- Prusky, D. and Plumbley, R.A., 1992. Quiescent infections of *Colletotrichum* in tropical and subtropical fruits. Pp 289-307. In *Colletotrichum: Biology, Pathology and Control* [edited by Bailey, J.A.; Jeger, M.J.] CAB International, Wallingford, UK.
- Quang, D. and Somsiri Sangchot, 2002. Infection of *Colletotrichum gloeosporioides* on Mango Fruit and its Resistance. *Journal. Agricultural. Science*, 33: 68-70
- Rojas, E.I., Rehner S.A., Samuels G.J., Van Bael S.A., Herre E.A., 2010. *Colletotrichum gloeosporioides* associated with *Theobroma cacao* and other plants in Panama: multilocus phylogenies distinguish pathogen and endophyte clades. *Mycologia*, 102: 1318-1338.
- Rojas-Martínez R.I, Zavaleta-Mejía E, Nieto-Angel D, 2008. Virulence and genetic variation of isolates of *Colletotrichum gloeosporioides* (Penz.) Penz. And Sacc. on mango (*Mangifera indica* L.) *Rev. mex. Phytopathology*, 26: 21–6.
- Sanders, G. M. and Korsten, L., 2003. Comparison of cross inoculation potential of South African avocado and mango isolates of *Colletotrichum gloeosporioides*. *Microbiology Research*, 158, 143–150.
- SAS, INC., 2008. SAS Guide for Personal Computers. Version 9.2, SAS Institute, Cary, NC

- Sharma, P.N., Kaur M., Sharma O.P., Sharma P., Pathania A., 2005. Morphological, pathological and molecular variability in *Colletotrichum capsici*, the cause of fruit rot of chillies in the subtropical region of northwestern India *Journal Phytopathology*, 153: 232–7.
- Shelke, R.B., Ingle, S.T., Patil, S.R., and Sonkar, U.V., 1997. Susceptibility to grape varieties to major disease in Vidarbha region. *Journal Soils and Crops*, 7: 150-152.
- Sutton, B.C., 1992. The genus *Glomerella* and its anamorph *Colletotrichum*. In: *Colletotrichum: Biology, Pathology and Control*. (eds. Bailey JA, Jeger MJ) Wallingford, UK: CAB International, 1-26.
- Syed, A. Hasan N, Rashida P, Syed A, Manzoor, H Muhammad I. Umar, M. Tayyab I. Fatima L, Taha M, Ahsan I., 2014. Evaluation of Various Mango Varieties against the Infection Dynamics of Powdery Mildew (*Oidium mangiferae* Bert.) *American Journal Plant Science*, 5: 2372-2377
- Thakur, R.P. and Rao, V.P., 1997. Variation in virulence and aggressiveness among pathology of *Sclerospora graminicola* on pearl millet. *Indian Phytopathology*, 50: 40-47.
- Than, P.P, Jeewon R, Hyde K.D, Pongsupasamit S, Mongkolporn O, Taylor P.W.J., 2008a. Characterization and pathogenicity of *Colletotrichum* species associated with anthracnose on chili (*Capsicum* species) in Thailand. *Plant Pathology*, 57: 562–572.
- Than, P.P, Shivas R.G, Jeewon R, Pongsupasamit S, Marney T.S, 2008b. Epitypification and phylogeny of *Colletotrichum acutatum* J.H. Simmonds. *Fungal Diversity*, 28: 97–108.



© The Author(s)

This is an  Open Access article licensed under a Creative Commons license: Attribution 4.0 International (CC-BY).